



United States Department of Agriculture

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen



Non-Discrimination Statement

Non-Discrimination Policy

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers, employees, and applicants for employment on the bases of race, color, national origin, age, disability, sex, gender identity, religion, reprisal, and where applicable, political beliefs, marital status, familial or parental status, sexual orientation, or all or part of an individual's income is derived from any public assistance program, or protected genetic information in employment or in any program or activity conducted or funded by the Department. (Not all prohibited bases will apply to all programs and/or employment activities.)

To File an Employment Complaint

If you wish to file an employment complaint, you must contact your agency's [EEO Counselor](#) (PDF) within 45 days of the date of the alleged discriminatory act, event, or in the case of a personnel action. Additional information can be found online at http://www.ascr.usda.gov/complaint_filing_file.html.

To File a Program Complaint

If you wish to file a Civil Rights program complaint of discrimination, complete the [USDA Program Discrimination Complaint Form](#) (PDF), found online at http://www.ascr.usda.gov/complaint_filing_cust.html, or at any USDA office, or call (866) 632-9992 to request the form. You may also write a letter containing all of the information requested in the form. Send your completed complaint form or letter to us by mail at U.S. Department of Agriculture, Director, Office of Adjudication, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, by fax (202) 690-7442 or email at program.intake@usda.gov.

Persons with Disabilities

Individuals who are deaf, hard of hearing or have speech disabilities and you wish to file either an EEO or program complaint please contact USDA through the Federal Relay Service at (800) 877-8339 or (800) 845-6136 (in Spanish).

Persons with disabilities who wish to file a program complaint, please see information above on how to contact us by mail directly or by email. If you require alternative means of communication for program information (e.g., Braille, large print, audiotape, etc.) please contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

Supplemental Nutrition Assistance Program

For any other information dealing with Supplemental Nutrition Assistance Program (SNAP) issues, persons should either contact the USDA SNAP Hotline Number at (800) 221-5689, which is also in Spanish or call the [State Information/Hotline Numbers](#).

All Other Inquiries

For any other information not pertaining to civil rights, please refer to the listing of the [USDA Agencies and Offices](#) for specific agency information.

Table of Contents

Executive Summary	1
Introduction	2
Development of the Pollinator Attractiveness Crop List	5
Use of the Pollinator Attractiveness Crop List	8
Table 1. Summary of the attractiveness to <i>Apis</i> and non- <i>Apis</i> bees of crops grown in the U.S., whether crop requires bee pollination and if so, whether managed pollinators are used.	9
Table 2. Additional crops identified in the 40 CFR crop groupings and their attractiveness to <i>Apis</i> and non- <i>Apis</i> bees, whether crop requires bee pollination and if so, whether managed pollinators are used.....	22
Table 3. List of references cited in Tables 1 and 2	40

Executive Summary

The protection of bee pollinators, *e.g.*, honey bees (*Apis mellifera*) and non-*Apis* bees in the United States has grown increasingly important, because declines in their populations have the potential to impact food security due to loss of pollination services. Several key factors have been implicated in overall honey bee colony losses, including pesticides. The U.S. Environmental Protection Agency (EPA) risk assessment process for pesticides includes an evaluation of risk to bee pollinators. This document provides a compilation of information on the attractiveness of crops grown in the United States to pollinating bees as food sources of pollen and nectar, and agronomic practices that are relevant to the interactions between these insects and the crops. The information provides a starting point for the risk assessment process for pollinating bees in terms of determining the potential for exposure to pesticide applications on these crops. In addition, the information contained in this resource will help to inform decisions to pursue further refinements in the risk assessment process as well as options for risk mitigation.

Introduction

The number of managed honey bee colonies in the U.S. has declined over the past 60 years and there are indications that the number and diversity of insect pollinators in general have also declined in North America¹. Additionally, populations of some non-*Apis* bee species have declined in recent decades^{2,3,4}. Although multiple factors have been associated with declines in *Apis* and non-*Apis* bees (e.g., arthropod pests, disease, poor nutrition, decreases in the diversity of food resources for bees, loss of habitat, lack of genetic diversity and pesticides), no single factor or specific combination of factors has been identified as the principal cause⁵. The U.S. Department of Agriculture (USDA) has been tasked by Congress to identify and to develop means to mitigate the causes of honey bee declines⁶. Although pesticides alone have not been implicated as the principal cause of overall bee pollinator declines, the EPA and USDA have been working collaboratively to understand the potential role that pesticides may be playing, particularly in combination with other identified factors.

In September, 2012, scientists from the EPA Office of Pesticide Program's Environmental Fate and Effects Division, in collaboration with Health Canada's Pest Management Regulatory Agency and the California Department of Pesticide Regulation, presented a White Paper⁷ to the Federal Insecticide, Fungicide and Rodenticide Act Scientific Advisory Panel (SAP) on a proposed framework for assessing risks of pesticides to bees in order to protect pollination services, production of hive products, and bee pollinator biodiversity. The proposed process for

¹ NAS. 2007. Status of Pollinators in North America. National Research Council of the National Academies. The National Academies Press, Washington DC. ISBN 978-0-309-10289-6.

² Grixti, J.C., L.T. Wong, S.A. Cameron, and C. Favret. 2009. [Decline of bumble bees \(*Bombus*\) in the North American Midwest](#). Biological Conservation 142:75–84.

³ Cameron, S.A., J.D. Lozier, J.P. Strange, J.B. Koch, N. Cordes, L.F. Solter, and T.L. Griswold. 2011. [Patterns of widespread decline in North American bumble bees](#). Proceedings of the National Academy of Sciences 108:662-667.

⁴ Bartomeus, I., J.S. Ascher, J. Gibbs, B.N. Danforth, D.L. Wagner, S.M. Hedtke, and R. Winfree. 2013. Historical changes in northeastern US bee pollinators related to shared ecological traits. Proceedings of the National Academy of Sciences 110(12): 4656 – 4660 doi: 10.1073/pnas.1218503110. <http://www.pnas.org/content/110/12/4656.short>

⁵ USDA. 2013. Report on the National Stakeholders Conference on Honey Bee Health. National Honey Bee Stakeholder Conference Steering Committee. Alexandria, Virginia. October 15 – 17, 2012. <http://www.usda.gov/documents/ReportHoneyBeeHealth.pdf>

⁶ USDA. 2007. Colony Collapse Disorder Action Plan, CCD Steering Committee, June 20, 2007. http://www.ars.usda.gov/is/br/ccd/ccd_actionplan.pdf

⁷ <http://www.epa.gov/scipoly/sap/meetings/2012/091112meeting.html>

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

assessing risks to bees, using honey bees as a surrogate for non-*Apis* bees as well, serves as a means to advance the science and to allow EPA to quantitatively assess the potential risk of pesticides to managed bees (*Apis* and non-*Apis*) and feral non-*Apis* bees.

The proposed framework for assessing risk to bees is similar to the process used by the EPA Office of Pesticide Programs for other taxa⁸; however, the biology of insect pollinators such as honey bees presents special considerations, which warrant a dedicated framework. Bee species can exhibit a wide range of social interaction/structure, with social structure of the various species affecting routes for potential exposure to pesticides. Often, the major commercial bee pollinators are either honey bees or bumble bees, but in some cases certain solitary bee species are also used commercially, e.g., blue orchard bees (*Osmia lignaria*) and alfalfa leafcutter bees (*Megachile rotundata*).

It is also important to note that risk assessors may have to evaluate a wide variety of plant types as a result of pesticide use patterns, from forestry and ornamental uses to use in crops, such as corn (*Zea mays*) and canola (*Brassica napus*). These uses may differ in regards to need and timing of commercial insect pollination services. However, some crops may be pollinator-dependent or pollinator-attractive when in bloom, but they may be typically harvested before flowering (e.g., lettuce) and would not represent a route of exposure based on typical cultivation practices. In other cases, some crop flowers are visited by solitary bees or bumble bees but not by honey bees. All of these pieces of information are essential to the understanding of bee pollinator visitation to a plant/crop of interest and the consequent need to assess the risk to bee pollinators from a pesticide application to this plant/crop.

As the SAP highlighted in the following conceptual model for the tiered approach in risk assessments to *Apis* and non-*Apis* bee pollinators (**Figure 1, Boxes 2a and 2b**), the first step of a risk assessment is to evaluate whether there is the potential for exposure. Therefore, information on the pollination biology of each plant/crop is needed to determine if bees are likely to visit the plants that are identified for pesticide applications. In addition, a risk assessor also needs information on the application rate, timing, method of application, and environmental fate of a pesticide in order to evaluate potential routes of exposure. Together, these pieces of information enable a risk assessor to determine if the proposed application of a pesticide leads to probable routes of exposure that could coincide with the timing of bee pollinator visitation to the plant/crop that is under consideration. In all cases, registrants provide information on the

⁸ USEPA. 2004. Overview of the Ecological Risk Assessment Process in the Office of Pesticide Programs. U.S. Environmental Protection Agency, Endangered and Threatened Species Effects Determinations. <http://www.epa.gov/espp/consultation/ecorisk-overview.pdf> . Last accessed 1/6/15.

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

application rate, timing, method of application, and environmental fate of a chemical when a new pesticide use pattern is proposed. However, to complement the use information, a comprehensive and robust source of information on the pollination ecology of the plants to which the pesticide is to be applied (*i.e.*, target plant) is needed to evaluate whether proposed uses for pesticide applications represent a potential exposure to either adult or immature (brood) bees.




Figure 1. Example tiered approach for assessing risk to honey bees from foliar spray applications.

Development of the Pollinator Attractiveness Crop List

In order to assess the potential for *Apis* and non-*Apis* bees to be exposed to pesticides applied to various crops, relevant data and agronomic practices connected to the pollination of these plants/crops are needed. These data include a measure of attractiveness to pollinating bees (*i.e.*, honey bees, bumble bees, and solitary bees) to each plant and/or crop, the phenology of the bloom period, use/non-use of managed pollinators, and the acreage of the various crop/plant groups in the United States.

The goal of this effort is to compile information on the attractiveness of crops grown in the United States to pollinating *Apis* and non-*Apis* bees as food sources of pollen and nectar. To achieve this goal, the EPA, USDA, and Michigan State University Extension Entomologist

Dr. Rufus Isaacs initiated a project to gather the relevant information to serve as a resource for pesticide risk assessments. The tables described here entitled “*Bee Pollinator Attractive Crops List*” (**Tables 1** and **2**) were developed to provide a relative rating of the degree to which honey bees, bumble bees, and solitary bees utilize the various crops grown in the United States. This list was informed by previous work conducted and recently published by the European Food Safety Authority (EFSA)⁹ for assessing pesticide risks to bees and adapted for the specific situations and regulatory data needs for the EPA.

Information for the Bee Pollinator Attractive Crops List (**Tables 1** and **2**) was collected from multiple sources, including peer-reviewed published information, university and agricultural extension resources, and expert opinion based on experience with the pollination of specific crops. The published information included the key texts of McGregor¹⁰, Free¹¹, and Delaplane and Mayer¹². Additionally, primary research publications were used where appropriate, and these are listed in **Table 3**. Online or published articles from expert knowledge of specific agricultural crop systems were also used to complete these entries, based on the experience of entomologists that work in crop pollination and bee keeping or from state agricultural extension

⁹ European Food Safety Authority, 2013. EFSA Guidance Document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees). Appendix D. EFSA Journal 2013; 11(7):3295, 266 pp., doi: 10.2903/j.efsa.2013.3295.

¹⁰ McGregor SE, 1976. Insect pollination of cultivated crop plants. Agricultural Handbook No. 496. Ed USDA Agricultural Research Service W, D.C, USA.

¹¹ Free JB. 1993. Insect Pollination of crops, 2nd edn. Academic Press: London, UK.

¹² Delaplane, K. S. & Mayer, D. F. (2000). Crop Pollination by Bees. – New York, Oxon (CABI Publishing).

agencies. When expert opinion was used as the source of information, the source is also identified in **Table 3**. If information could not be identified from publications or expert opinion to address a specific aspect of the pollination biology of the plant or its associated typical agronomic practice, the corresponding cells in **Tables 1** and **2** were left blank and this aspect of the crop remains an uncertainty.

There are specific considerations for some of the data listed in **Tables 1** and **2** related to the rating of attractiveness of pollen and/or nectar resources to bees. For each of the crops listed, the degree to which pollen and nectar are attractive and used by honey bees is listed using a scale where "-" = not attractive, "+" = attractive under certain conditions, and "++" = high attractiveness in all cases. The same rating system is used for bumble bees and for solitary bees with the major groups of solitary bees likely to be found at flowers of each crop listed, if known. The ratings for bumble bee and solitary bee taxa do not address pollen and nectar separately, however. The different attractiveness ratings are based on the degree to which information qualitatively indicates that they are used by bees. If the cited information indicates that certain bees frequently visit and extensively use a particular floral resource, then it is given the classification of "++" for the respective type of bee. If, however, information indicates that certain bees only visit a crop infrequently (e.g., only under conditions of few alternative food sources) or few bees are noted to forage on a given crop resource, it is given the classification of "+" for the appropriate taxa of bee. Despite the relatively lower level of attractiveness compared to crops with a "++" rating, it is important to note that crops designated with a "+" may still become a major source of food for bees depending on the environmental conditions. For example, a crop that under normal conditions bees would only minimally use as a forage source, could be extensively used during certain time periods due to the lack of alternative available forage (e.g., drought, flooding, etc.). Additionally, nearby competing crops which may be more pollinator attractive may draw away some groups of pollinators due to the ease of obtaining pollen/nectar. Finally, when the various groups of bees are noted to be absent from a particular crop or resource, this crop is noted with a "-" for the appropriate type of bee. When crop specific information was available, attractiveness ratings are based on the inherent attractiveness of the crop to pollinating bees and not based on specific agronomic practices such as harvest prior to bloom. It is assumed that a crop that is harvested prior to bloom would be "unattractive" to pollinating bees as it would not provide flowers for visitation during typical cultivation.

There are also considerations specifically related to **Table 1** given that for most of the crops data were already available in the EFSA guidance document¹³. The first consideration is that if the specific attractiveness rating was not listed in the EFSA guidance document for solitary

¹³ *Ibid.* European Food Safety Authority, 2013.

bees, but rather only the type of bee was listed to denote floral visitation by that bee, then a “+” rating is applied in **Table 1** to denote that solitary bees visit the specific crop. The second consideration is that in many cases data were already available for the various crops in the EFSA document. Where additional data could not be found, the data in **Table 1** show the attractiveness ratings simply based on the EFSA guidance document. If other data provided clarification on additional bee taxa not addressed by EFSA and/or simply confirmed the EFSA data, then the EFSA data were combined with the additional data source noted in the reference. In some cases, other data sources modified the ratings provided by EFSA and these citations are provided in the references column.

In some cases, information is lacking for a crop, but another crop is identified as a surrogate for the rating of attractiveness to bees given similarity in plant family or crop group. In these cases, the surrogate is identified in the “Reference” column and serves as a reasonable assumption of attractiveness to bees. It should be noted, however, that there is uncertainty in the use of surrogate plants for the rating, as attractiveness may vary even within plant families.

Whether a crop requires bee pollination or not includes specific considerations related to the agronomic practices of the crop. The entry for 'requires bee pollination' refers to harvestable, productive crop yields rather than any specific level of fruit set. Consequently, if a crop “requires bee pollination” then the specific crop requires either specific bee taxa or bees in general to produce productive and harvestable crop yields. If a crop does not require bee pollination, then the specific crop attains harvestable and productive crop yields via other pollination methods aside from bees, such as through wind, mammal, or other invertebrate (e.g., butterfly) pollination. Whether the crop requires bee pollination is listed based on the information provided in the cited reference, which provides information on whether the crop has a dependence on bee-mediated pollen transfer for the production of seeds, nuts, or fruit. This information is specifically for the production of the edible crop parts that will be harvested and sold. Many crops do not require bees for pollination to produce marketable yields, yet they do require bees for breeding or seed production (*i.e.*, intended for crop propagation), which is typically a very small proportion of the total crop acreage. In these cases the crops are stated as requiring bee pollination and the “Notes” column of **Tables 1 and 2** state if the requirement is only for seed production. Finally, some crops may produce flowers during the typical production of the harvestable part of the crop but does not require bee pollination. In these cases, exposure to bees is assumed and the attractiveness ratings of the crop to bees are specified in **Tables 1**

and **2**.

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Many crops in **Tables 1** and **2** employ commercial honey bee colonies (or colonies of other types of bees) that are rented by the grower and provided by a beekeeper to provide pollination services. If that is likely to happen within a cropping system within the United States, based on the information provided in the cited reference, it is listed as "Yes" in the "Uses Managed Pollinators" column. If that does not happen based on the designation of not requiring bee pollination, then the column has a "No" entry based on inference drawn from the column on bee pollination requirements.

Use of the Pollinator Attractiveness Crop List

As part of its pesticide ecological risk assessments, the EPA intends to consider the information on pollinator attractiveness provided in **Tables 1** and **2** in determining the potential for bees to be exposed to pesticides from the crop itself following application to a specific crop. If a risk assessment is warranted, other information included in **Tables 1** and **2** can assist in refining the scope of the risk assessment. The other information includes the spatial extent of treated crops, the timing of application in relation to likely periods of bee visitation, and agronomic practices that may affect the exposure of the bees (*e.g.*, harvesting prior to bloom). Where necessary, information in this list may be supplemented with additional information on a case-by-case basis (*e.g.*, crop and region-specific information from local agricultural extension experts).

Table 1. Summary of the attractiveness to *Apis* and non-*Apis* bees of crops grown in the U.S., whether crop requires bee pollination and if so, whether managed pollinators are used.

Also summarized is the bearing acreage of the crop, the extent to which the crop is used in seed production and whether the crop is harvested prior to bloom. The degree to which pollen and nectar are attractive is listed using a scale where "-" = not attractive, "+" = attractive under certain conditions, and "++" = high attractiveness; entry "N/AV" specifies when crop-specific data are unavailable; entry "N/AP" specifies when crop-specific data are not applicable.

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Alfalfa	<i>Medicago sativa</i>	+	++	+	++ Alfalfa leafcutting bee, Alkali bee	Yes	Yes	1	17,763,000	2011: 6600 acres	Yes	Also grown for seed on a small % of acreage where both Megachilidae and <i>Apis</i> are contracted; alfalfa grown for forage harvested at 10% bloom.
Almonds	<i>Prunus amygdalus</i> ; <i>P. communis</i> ; <i>Amygdalus communis</i>	++	+	+	+ <i>Osmia</i>	Yes	Yes	1	780,000		No	
Anise, badian, fennel, corian, juniper berries	anise (<i>Pimpinella anisum</i>); badian or star anise (<i>Illicium verum</i>); caraway (<i>Carum carvi</i>); coriander (<i>Coriandrum sativum</i>); cumin (<i>Cuminum cyminum</i>); fennel (<i>Foeniculum vulgare</i>); juniper berries (<i>Juniperus communis</i>)	+	+	+	+	Yes (not juniper berries)	No	2	N/AV		No	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Apples	<i>Malus pumila</i> ; <i>M. sylvestris</i> ; <i>M. communis</i> ; <i>Pyrus malus</i>	++	+	+	++ <i>Andrena</i> , <i>Anthidium</i> , <i>Halictus</i> , <i>Osmia</i> , <i>Anthophora</i> , <i>Habropoda</i>	Yes	Yes	1	327,800		No	
Apricots	<i>Prunus armeniaca</i>	++	++	++	+ <i>Osmia</i>	Yes	Yes	3	12,150		No	
Artichokes	<i>Cynara scolymus</i>	+	+	+	+	Yes	No	3,4, 81	7,000		Yes	
Asparagus	<i>Asparagus officinalis</i>	+	+	N/AV	N/AV	Yes	Yes	1	24,500		Yes	Requires pollination only when grown for seed; small % of acreage
Avocados	<i>Persea americana</i>	+	+	N/AV	+	Yes	Yes	1	59,950		No	
Bananas	<i>Musa sapientum</i> ; <i>M. cavendishii</i> ; <i>M. nana</i>	-	+	-	-	No	No	5	1,000		No	
Barley	<i>Hordeum</i> spp.	-	-	-	-	No	No	3	3,000,000		No	Wind pollinated
Beans	<i>Phaseolus</i> spp.	+	+	+	N/AV	No	No	3	77,200		No	Acreage is for snapbeans
Blueberries	fruits of the genus <i>Vaccinium</i>	+	+	++	++ <i>Andrena</i> , <i>Colletes</i> , <i>Osmia</i> , <i>Anthophora</i> , <i>Xylocopa</i>	Yes	Yes	1	77,700		No	Acreage is only for cultivated blueberries; <i>Apis M.</i> and Megachilidae used in commercial pollination.
Broad beans, horse beans, dry	<i>Vicia faba</i>	++	++	++	+ <i>Anthophora</i> , <i>Eucera</i> , <i>Megachile</i> , <i>Xylocopa</i>	Yes		5	1,311,300		No	
Buckwheat	<i>Fagopyrum esculentum</i>	+	++	+	+	Yes	Yes	5, 73	33,678		No	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Cabbages and other brassica	Chinese, mustard cabbage, pak-choi (<i>Brassica chinensis</i>); white, red, Savoy cabbage, Brussels sprouts, collards, kale and kohlrabi (<i>Brassica oleracea</i> all varieties except <i>botrytis</i>)	++	++	+	+	Yes	Yes	1	Cabbage 60,180 (Annual); Brussels sprouts 7,569 (Census); Kale 6,256 (Census); Collards 12,542 (Census)		Yes	Requires pollination only when grown for seed. Small % of acreage
Carobs	<i>Ceratonia siliqua</i> , Carob tree, locust bean	+	+	+	+	Yes	No	49, 74				Flowers visited mainly by flies and wasps
Carrots	<i>Daucus carota</i>	+	+	+	+ <i>Megachile rotundata</i>	Yes	Yes	1, 3	71,400 Fresh Market; 13,310 Processing	2012: 4941 acres	Yes	Requires pollination only when grown for seed. Small % of acreage
Castor oil seed	<i>Ricinus communis</i>	+	-	N/AV	N/AV			EFSA	N/AV	Yes	No	
Cauliflowers and broccoli	<i>Brassica oleracea</i> var. <i>botrytis</i> , subvarieties <i>cauliflora</i> and <i>cymosa</i> , includes headed broccoli	++	++	+	+ <i>Andrenidae</i> , <i>Nomadidae</i> , <i>Megachilidae</i>	Yes	Yes	5	163,730 Fresh market and processing		Yes	Requires pollination only when grown for seed; small % of acreage
Cherries	Mazzard, sweet cherry (<i>Prunus avium</i> ; <i>Cerasus avium</i>); hard-fleshed cherry (var. <i>duracina</i>); heart cherry (var. <i>juliana</i>)	++	+	+	++ <i>Osmia</i>	Yes	Yes	1	86,790 Sweet; 36,500 Tart		No	
Chestnuts	<i>Castanea</i> spp.: <i>C. vesca</i> ; <i>C. vulgaris</i> ; <i>C. sativa</i> .	++	++	+	+	Yes	Yes	3	3,784			

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Chick peas	Chickpea, Bengal gram, garbanzos (<i>Cicer arietinum</i>)	+	++	+	+ <i>Osmia</i> , <i>Megachile</i>	No	No	72	213,600; Note: Included in All Dry Bean Acres			Self-pollinated
Chicory roots	<i>Cichorium intybus</i> subsp. <i>sativum</i>	+	+	N/AV	+ <i>Andrena</i> , <i>Anthidium</i> , <i>Halictus</i> , <i>Osmia</i>	Yes	N/AV	EFSA, 3	N/AV		Yes	
Chillies and peppers	Red and cayenne pepper, paprika, chillies (<i>Capsicum frutescens</i> ; <i>C. annuum</i>); allspice, Jamaica pepper (<i>Pimenta officinalis</i>)	+	-	++	+	Yes	No	1	71,200 Chile and Bell			May be grown in glasshouses, with bumble bees for pollination
Clover for forage and silage	<i>Trifolium</i> spp. Various species grown for pasture, green fodder or silage	++	++	+	++ <i>Megachile</i> , <i>Osmia</i> , <i>Andrena</i> , <i>Anthidium</i>	Yes	Yes	1,5, 89, 102, 103	28,506 White, Red and Crimson		Yes	Requires pollination only when grown for seed; small % of acreage
Coffee, green	<i>Coffea</i> spp. (<i>arabica</i> , <i>robusta</i> , <i>liberica</i>)	+	-	N/AV	+	Yes	No	5	7300	Yes	No	Acreage related to all coffee, not specific to green coffee
Corn	<i>Zea mays</i>	+	-	+	+	No	No	3	87,668,000			Wind pollinated, but can be visited during pollen shedding

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Cotton	Upland cotton (<i>Gossypium hirsutum</i>) Pima Cotton (<i>Gossypium barbadense</i>)	-	+	+	<i>Halictus, Anthophora, Xylocopa, Megachile, Nomia, Ptilothrix</i>	No	No	5, 104, 105, 106, 107, 108, 109, 110, 111	7,664,400	Historical use of bees for hybrid seed production; however, hybrid cotton seed production is no longer considered economically viable		Used by some beekeepers for honey production
Cow peas	Cowpea, blackeye pea/bean (<i>Vigna unguiculata</i>)	-	+ ³	+	+	Yes	N/AV	11	39,100 Blackeye Peas, Included with All Dry Beans		No	
Cranberries	American cranberry (<i>Vaccinium macrocarpon</i>)	+	+	++	++ <i>Andrena, Agapostemon, Melitta, Megachile</i>	Yes	Yes	1	40,300		No	
Cucumbers and gherkins	<i>Cucumis sativus</i>	+	+	+	+ <i>Melissodes Andrena</i>	Yes	Yes	1	40,060 Fresh; 82,100 for Pickles	Yes		Small seed acreage
Currants	Black (<i>Ribes nigrum</i>); red and white (<i>R. rubrum</i>)	-	+	++	+ <i>Anthophora</i>	Yes	No	5	580 Total		No	
Dates	<i>Phoenix dactylifera</i>	+	+	N/AV	N/AV	No	No	3	8,400		No	Wind pollinated
Eggplants	<i>Solanum melongena</i>	-	-	++	+	Yes	No	5	5,004		No	Requires pollination only when grown for seed; small % of acreage
Elder	<i>Sambucus nigra</i>	+	+	+	+	No	No	6	N/AV			
Figs	<i>Ficus carica</i>	-	-	-	-	No	No	5	8,600		No	Wasp pollinated

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Garlic	<i>Allium sativum</i>	+	+	N/AV	+ <i>Halictus</i> , <i>Osmia</i>	Yes	No	3	23,900		Yes	Requires pollination only when grown for seed; small % of acreage
Gooseberry	<i>Ribes grossularia</i>	-	+	++	+	Yes	No	5	N/AV		No	Little production in US
Grapefruit (inc. pomelos)	<i>Citrus maxima</i> ; <i>C. grandis</i> ; <i>C. paradisi</i>	++	++	+	N/AV	No	No	3, 9	73,300 (no pomelos)		No	
Grapes	<i>Vitis vinifera</i>	+	-	-	-	No	No	5	962,100		No	Wind pollinated
Grasses for forage; Sil	Including inter alia: bent, redtop, fiorin grass (<i>Agrostis</i> spp.); bluegrass (<i>Poa</i> spp.); Columbus grass (<i>Sorghum almum</i>); fescue (<i>Festuca</i> spp.); Napier, elephant grass (<i>Pennisetum purpureum</i>); orchard grass (<i>Dactylis glomerata</i>); Rhodes grass (<i>Chloris gayana</i>); <i>Phleum</i> , <i>Agropyron</i> , <i>Elymus</i> , <i>Phalaris</i> , <i>Koeleria</i> , <i>Stipa</i> , <i>Danthonia</i> , <i>Deschampsia</i> , <i>Bromus</i> , <i>Trisetum</i> , <i>Calamagrostis</i> , <i>Carex</i> and <i>Juncus</i>	+	-	-	-	No	No	5	35,328,000		Yes	Wind pollinated, source of pollen only when no other forage sources are available
Groundnuts, with shell, peanuts	<i>Arachis hypogaea</i>	+	N/AV	+	+ <i>Lasioglossum</i> , <i>Megachile</i> , <i>Anthidium</i> , <i>Nomia</i>	N/AV	N/AV	EFSA	1,042,000			

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Hazelnuts, with shell (filberts)	<i>Corylus avellana</i>	+	-	-	-	No	No	50	29,000			
Hemp	<i>Cannabis sativa</i>	+	-	+	N/AV	No	No	51	N/AV			Wind pollinated
Hops	<i>Humulus lupulus</i>	+	-	-	-	No	No	7, 82	35,224			
Kiwi fruit	<i>Actinidia chinensis</i>	+	+	+	+	Yes	Yes	1	4,200			
Leeks, other alliaceous vegetables	Leeks (<i>Allium porrum</i>); chives (<i>A. schoenoprasum</i>); other alliac	+	++	+	+ <i>Osmia</i> , <i>Hoplitis</i>	Yes	No	3, 5	N/AV		Yes	Requires pollination only when grown for seed; small % of acreage
Leguminous for silage	Including inter alia: birdsfoot trefoil (<i>Lotus corniculatus</i>); lespedeza (<i>Lespedeza</i> spp.); kudzu (<i>Pueraria lobata</i>); sesbania (<i>Sesbania</i> spp.); sainfoin, esparcette (<i>Onobrychis sativa</i>); sulla (<i>Hedysarum coronarium</i>)	+	++	++	++ <i>Anthidium</i> , <i>Anthophora</i> , <i>Lasioglossum</i> , <i>Megachile</i> , <i>Osmia</i> , <i>Xylocopa</i>	Yes	Yes	3, 8, 102, 103	Birdsfoot - Not Published; 3,219 Lespedeza			Trefoil is valuable honey plant for beekeepers. Potential use of the <i>Megachilidae</i> to pollinate sweet clover and sanfoin
Leguminous vegetables	<i>Vicia faba</i>	++	++	++	+ <i>Anthophora</i> , <i>Eucra</i> , <i>Megachile</i>	Yes	No	1	N/AV		No	
Lemons/ limes	Lemon (<i>Citrus limon</i>); sour lime (<i>C. aurantifolia</i>); sweet lime (<i>C. limetta</i>)	++	++	N/AV	+	No	No	5	55,000 Lemons (Annual) 820 Limes (Census)			
Lentils	<i>Lens esculenta</i> ; <i>Ervum lens</i>	+	+ ³	-	+ <i>Megachile</i>	No	No	52	347,000			

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Lettuce	<i>Lactuca sativa</i>	+	+	+	+	No	No	3, 5	259,100 Head, Leaf and Romaine		Yes	Self-pollinating
Linseed	<i>Linum usitatissimum</i> Flaxseed.	-	-	-	-	No	Yes	90	N/AV			Extensively grown in the Dakotas and the Canadian Prairies.
Lupins	<i>Lupinus alba</i> , <i>L. angustifolia</i> , <i>L. luteus</i> .	+	-	++	+	Yes	No	91	N/AV			Requires pollination only when grown for seed; small % of acreage
Melonseed	<i>Cucumis melo</i> , includes seeds of other Cucurbitaceae	+	+	+	+ <i>Ceratina</i> , <i>Peponapis</i> , <i>Melissodes</i> , <i>Agapostemon</i>	Yes	Yes	5	N/AV		No	
Mushrooms and truffles	Edible mushrooms	N/AP	N/AP	N/AP	N/AP	No	No	40				Produced indoors in the dark, no bee pollination required
Mustard seed	White mustard (<i>Brassica alba</i> ; <i>B. hirta</i> ; <i>Sinapis alba</i>); black mustard (<i>Brassica nigra</i> ; <i>Sinapis nigra</i>) <i>Brassica juncea</i>	++	++	+	+	Yes	N/AV	5		43,400	No	<i>B. juncea</i> extensively grown on Great Plains and southern Canadian prairies; is 2/3 self fertile and 1/3 out crossing, so bees partially required
Oat	<i>Avena</i> spp., mainly <i>Avena sativa</i>	-	-	-	-	No	No	3	1,030,000			Wind pollinated
Okra	<i>Abelmoschus esculentus</i> ; <i>Hibiscus esculentus</i>	+	+	+	+	Yes	No	5	2,377			
Olives	<i>Olea europaea</i>	+	-	N/AV	N/AV	No	No	3	44,000			

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Onions	<i>Allium cepa</i>	+	+	-	+ <i>Halictus</i> , <i>Nomia</i>	Yes	Yes	5	143,340	dry bulb: 2010: 73213 acres	Yes	Requires pollination only when grown for seed; small % of acreage
Oranges	Common, sweet orange (<i>Citrus sinensis</i>); bitter orange (<i>C. aurantium</i>)	++	++	+	+ <i>Andrena</i> , <i>Xylocopa</i>	No	Yes	9	613,000			Variable among orange cultivars; honey bees brought to groves for orange blossom honey
Peaches/ nectarines	<i>Prunus persica</i> ; <i>Amygdalus persica</i> ; <i>Persica laevis</i>	+	+	+	+ <i>Osmia</i>	Yes	Yes	1	112,880 Peaches; 26,400 Nectarines			
Pears	<i>Pyrus communis</i>	+	+	+	+ <i>Osmia</i> , <i>Andrena</i>	Yes	Yes	1	54,400			
Peas	Garden pea (<i>Pisum sativum</i>); field pea (<i>P. arvense</i>)	+	+	+	+ <i>Eucera</i> , <i>Xylocopa</i>	No	No	7	797,000	2013; 406 acres		
Peppermint	<i>Mentha</i> spp.: <i>M. piperita</i>	+	++	++	+	No	No	39	68,800			Peppermint oil is produced from vegetative growth, without flowering or seed production
Persimmons	<i>Diospyros kaki</i> ; <i>D. virginiana</i>	+	+	+	+	Yes	No	5	4,968			
Pistachios	<i>Pistacia vera</i>	-	-	-	-	No	No	53	178,000			Wind pollinated
Plums and sloes	Greengage, mirabelle, damson (<i>Prunus domestica</i>); sloe (<i>P. spinosa</i>)	+	+	+	+ <i>Osmia</i> , <i>Anthophora</i>	Yes	Yes	1	82,780			

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Poppy seed	<i>Papaver somniferum</i>	++	-	N/AV	N/AV	No	N/AV	EFSA, 92	N/AV			Mainly self fertile although cross pollination via insect, bees does occur
Potatoes	<i>Solanum tuberosum</i> Irish potato	-	-	+	+ <i>Andrena</i>	Yes	No	3	1,052,000			Requires pollination only for breeding; small % of acreage
Pumpkins, squash and gourds	<i>Cucurbita</i> spp., includes marrows	+	+	++	+ <i>Agapostemon</i> , <i>Melissodes</i> , <i>Peponapis</i>	Yes	Yes	5	91,700 Pumpkins and Squash			
Pyrethrum, dried	<i>Chrysanthemum cinerariifolium</i>	+	+	+	+	No	No	3, 81	N/AV			
Quinces	<i>Cydonia oblonga</i> ; <i>C. vulgaris</i> ; <i>C. japonica</i>	+	+	N/AV	N/AV	N/AV	N/AV	EFSA	N/AV			
Rapeseed (including canola)	<i>Brassica napus</i> var. <i>oleifera</i>	++	++	+	++ <i>Megachile</i>	Yes	Yes	1,3,5	1,264,500 Canola; 1,700 Rapeseed	2013: 1,500 acres		Managed bees needed for hybrid seed production
Raspberries	<i>Rubus idaeus</i>	+	+	++	+ <i>Osmia</i> , <i>Andrena</i> , <i>Coletes</i> , <i>Halictus</i>	Yes	Yes	1	17,300			
Rice, paddy	<i>Oryza</i> spp., mainly <i>Oryza sativa</i> .	-	-	-	-	No	No	3	2,468,000			Wind pollinated
Rye	<i>Secale cereale</i>	-	-	-	-	No	No	3	278,000			Wind pollinated
Rye grass for forage and silage	Italian ryegrass (<i>Lolium multiflorum</i>); English, perennial ryegrass (<i>L. perenne</i>).	-	-	-	-	No	No	3	N/AV			Wind pollinated

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Safflower seed	<i>Carthamus tinctorius</i>	+	+	N/AV	+	Yes	Yes	EFSA, 93	170,000			Safflower is basically self-pollinated, but bees or other insects are generally necessary for optimum fertilization and maximum yield
Serradella/ birdsfoot	<i>Ornithopus sativus</i>	+	++	N/AV	+ <i>Megachile</i>	Yes	N/AV	EFSA	N/AV			
Sesame seed	<i>Sesamum indicum</i>	+	++	N/AV	+	Yes	No	5	17,501			
Sorghum	<i>Sorghum bicolor</i> , spp. <i>bicolor</i>	+	-	N/AV	+	No	No	3, 83	6,910,000 Grain and Silage			
Soybeans	<i>Glycine soja</i>	+	+	+	+	No	No	1	75,869,000			
Spices	Including inter alia: bay leaves (<i>Laurus nobilis</i>); dill seed (<i>Anethum graveolens</i>); fenugreek seed (<i>Trigonella foenum-graecum</i>); saffron (<i>Crocus sativus</i>); thyme (<i>Thymus vulgaris</i>); turmeric (<i>Curcuma longa</i>)	+	+	+	+	No	No	5	N/AV			Attractiveness depends on the species
Spinach	<i>Spinacia oleracea</i>	-	-	-	-	No	N/AV	EFSA	31,440		Yes	
Strawberries	<i>Fragaria</i> spp.	+	+	+	+ <i>Andrena</i> , Halictids, <i>Osmia</i>	No	Yes	3	58,190			Not essential, but some growers add supplemental hives to compliment wind pollination

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Sugar beet	<i>Beta vulgaris</i> var. <i>altissima</i>	-	+	N/AV	+	Yes	No	3	1,154,200		Yes	Requires pollination only for breeding; small % of acreage
Sugar cane	<i>Saccharum officinarum</i>	-	-	-	-	No	No	3	905,600	2013: 907 acres		Wind pollinated
Sunflower seed	<i>Helianthus annuus</i>	++	++	++	++ <i>Halictus, Dieunomia, Megachile, Melissodes, Svastra, Xylocopa</i>	Yes	Yes	1	1,474,600	2013: 1,502,000 acres		
Sweet potatoes	<i>Ipomoea batatas</i>	+	+	+	+	Yes	No	5, 41, 78, 79	113,200			Propogated vegetatively, and requires pollination only for breeding. Small % of acreage
Tangerines, mandarins, clementines	Mandarin, tangerine (<i>Citrus reticulata</i>); clementine, satsuma (<i>C. unshiu</i>)	++	++	+	+ <i>Andrena, Xylocopa</i>	Yes	Yes	9	52100 Tangerines and Mandarins			
Tobacco ⁵	<i>Nicotiana tabacum</i>	+	-	+	+	No	No	44, 84	355,700		Yes	Typically deflowered as a standard production practice
Tomatoes	<i>Lycopersicon esculentum</i>	-	-	+	+	Yes	Yes	1	93,600 Fresh; 277,000 Processing			May be grown in glasshouses where bumble bees are needed for pollination
Triticale	<i>Triticum x Secale</i>	-	-	-	-	No	No	N/AV ⁶	61,428			Triticale is a cross between wheat (<i>Triticum</i>) and rye (<i>Secale</i>), both wind pollinated

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	Description	HB Poll. ¹	HB Nec. ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Ref No.	U.S. Bearing Acreage ²	Seed Production ⁷	Harvest Prior to Bloom	Notes
Turnips for fodder	<i>Brassica rapa</i> var. <i>rapifera</i> .	++	++	+	+	Yes	Yes	3	N/AV		Yes	Requires pollination only for breeding; small % of acreage
Vetches	Spring/common vetch (<i>Vicia sativa</i>).	++	+	++	++	Yes	No	42	3,441			
Viper's grass	<i>Scorzonera hispanica</i>	+	+	+	+	Yes	No	43	N/AV			*Note citation is not yet published*
Walnuts with shell	<i>Juglans</i> spp.: <i>J. regia</i> .	+	-	-	-	No	No	EFSA, 45	245,000			Wind pollinated
Watermelons	<i>Citrullus vulgaris</i>	+	+	+	+ <i>Agapostemon</i> , <i>Floridegus</i> , <i>Halictus</i> , <i>Hoplitis</i> , <i>Melissodes</i>	Yes	Yes	1	123,330			
Wheat	<i>Triticum</i> spp.: common (<i>T. aestivum</i>), durum (<i>T. durum</i>), spelt (<i>T. spelta</i>).	-	-	-	-	No	No	3	45,157,000			

† Major crops based on Appendix D in the EFSA bee risk assessment guidance document and their attractiveness to pollinating bees. The table also contains relevant agronomic information associated with each crop. The entry "N/AV" specifies when crop-specific data are unavailable. Where "EFSA" is listed as the reference for a specific crop in this table, the data from Appendix D in the EFSA bee risk assessment guidance are used as the sole source of information on attractiveness ratings as no additional data were identified.

¹ HB= honey bee; Pol = Pollen; Nec = Nectar

² Estimates from the Census of Agriculture have a 2012 harvested acreage date. NASS fruit estimates have a 2012 reference date and vegetables refer to 2013. Fruit estimates are in bearing acres. Field crops and specialty crops are reported in harvested acres. All Census estimates are reported in harvested acreage. N/AV = not available. Please refer to reference 48 in **Table 3** for the citation related to these data.

³ Extra-floral nectaries

⁴ Mainly on extra-floral nectaries

⁵ Unmanufactured tobacco

⁶ Extrapolation based on wheat and rye

⁷ Seed production refers to crops grown to produce seeds intended for crop propagation rather than for human or livestock consumption

Table 2. Additional crops identified in the 40 CFR crop groupings and their attractiveness to *Apis* and non-*Apis* bees, whether crop requires bee pollination and if so, whether managed pollinators are used.

The degree to which pollen and nectar are attractive is listed using a scale where "-" = not attractive, "+" = attractive under certain conditions, and "++" = high attractiveness; entry "N/AV" specifies when crop-specific data are unavailable. The table also contains relevant agronomic information associated with each crop.

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Arracha (PR) <i>Arracacia xanthorrhiza</i> (Apiaceae)	Roots and tuber vegetables	+	+	+	+	Yes	No	Extrapolated from carrot in Table 1	Bees important for seed production. Typically harvested prior to bloom.
Arrowroot <i>Maranta arundinacea</i> (Marantaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Chinese artichoke <i>Stachys affinis</i> (Lamiaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Jerusalem artichoke (Asteraceae)	Roots and tuber vegetables	+	+	+	+	No	No	38	Some genotypes produce viable seed which is generated by cross pollination by bees, mainly non- <i>Apis</i> .
Edible burdock (Asteraceae)	Roots and tuber vegetables	+	+	+	+	No	No	Extrapolated from Jerusalem artichoke above	
Edible canna (Cannaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Cassava (Euphorbiaceae)	Roots and tuber vegetables	-	-	-	-	No	No	3	
Turnip-rooted chervil (Apiaceae)	Roots and tuber vegetables	+	+	+	+	Yes	No	Extrapolated from coriander in Table 1	Requires pollination only when grown for seed. Typically harvested prior to bloom.
Chufa (Cyperaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Dasheen (Araceae)	Roots and tuber vegetables	+	+	N/AV	N/AV	No	No	46, 77	Cultivated primarily vegetatively
Ginger (PR) (Zingiberaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Ginseng (Araliaceae)	Roots and tuber vegetables	N/AV	N/AV	N/AV	+	No	No	75	
Horseradish (Brassicaceae)	Roots and tuber vegetables	+	+	+	+	No	No	Attractiveness extrapolated from radish below	Asexual reproduction through root propagation.
Leren (PR) (Marantaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Turnip rooted parsley (Apiaceae)	Roots and tuber vegetables	+	+	+	+	No	No	Extrapolated from parsley below	Bees important for seed production. Typically harvested prior to bloom.
Parsnip (Apiaceae)	Roots and tuber vegetables	+	+	+	+	Yes	No	3	Requires pollination only when grown for seed. Small % of acreage. Typically harvested prior to bloom.
Radish (Brassicaceae)	Roots and tuber vegetables	+	+	+	+ <i>Megachile</i>	Yes	Yes	3	Requires pollination only when grown for seed. Honey bees are the primary pollinators. Typically harvested prior to bloom.
Rutabaga and turnip (Brassicaceae)	Roots and tuber vegetables	++	++	+	+	Yes	Yes	3	Requires pollination only when grown for seed. Small % of acreage. Typically harvested prior to bloom.
Salsify, (oyster plant) (Asteraceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Salsify, spanish (Asteraceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Skirret (Apiaceae)	Roots and tuber vegetables	+	+	+	+	Yes	No	Extrapolated from carrot on Table 1	Bees important for seed production. Typically harvested prior to bloom.
Tanier (Araceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Yam bean (Fabaceae)	Roots and tuber vegetables	+	+	+	+	No	No	Extrapolated from Bean (lupinus) below	
True yam (Dioscoreaceae)	Roots and tuber vegetables	<i>Uncertainty^a</i>							
Chive, Chinese (Liliaceae)	Bulb vegetables	+	++	+	+	Yes	No	Extrapolated from chive above	Requires pollination only when grown for seed; small % of acreage
Daylily, bulb (Liliaceae)	Bulb vegetables	-	-	-	-	No	No	12	Primarily moth and butterfly pollinated
Elegans hosta (Liliaceae)	Bulb vegetables	+	-	+	+	No	No	7	
Fritillaria (Liliaceae)	Bulb vegetables	+	+	+	+	No	No	13	
Garlic, great headed (Liliaceae)	Bulb vegetables	+	+	+	+	No	No	3, 81	Rarely grown for seed
Garlic, serpent (Liliaceae)	Bulb vegetables	+	+	+	+	No	No	Extrapolated from great headed garlic above	Rarely grown for seed
Kurrrat (Liliaceae)	Bulb vegetables	+	++	N/AV	+ <i>Osmia</i> , <i>Hoplitis</i>	Yes	No	Extrapolated from leek in Table 1: 3, 5	Typically harvested prior to bloom. Requires pollination only when grown for seed; small % of acreage
Lily (Liliaceae)	Bulb vegetables	-	-	-	-	No	No	57	Rarely grown for seed
Onion (various varieties except green onion) (Liliaceae)	Bulb vegetables	+	+	+	+	No	No	3, 14, 81, Attractiveness extrapolated from green onion in Table 1	Requires pollination when grown for seed; small % of acreage, but locally important (CA, AZ)

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes	
Shallot (Liliaceae)	Bulb vegetables	+	+	+	+	No	No	3, 14, 81, Attractiveness extrapolated from green onion in Table 1	Requires pollination when grown for seed; small % of acreage, but locally important (CA, AZ)	
Amaranth (Amaranthaceae)	Leafy Vegetables	+	+	+	+	Yes		94	Crop harvested prior to bloom	
Arugula (Brassicaceae)	Leafy Vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Crop is harvested prior to bloom when not grown for seed production.	
Cardoon (Asteraceae)	Leafy Vegetables	+	+	+	+	Yes	No	81, Attractiveness extrapolated from artichoke in Table 1	Crop is harvested prior to bloom when not grown for seed production.	
Celery (Apiaceae)	Leafy Vegetables	+	+	+	+	Yes	No	3, Attractiveness to wild bees extrapolated from parsley	Crop harvested prior to bloom. Bees important for seed production.	
Celtuce (Asteraceae)	Leafy Vegetables	+	+	+	+	No	No	Extrapolated from lettuce in Table 1	Crop is harvested prior to bloom when not grown for seed production.	
Chervil (Apiaceae)	Leafy Vegetables	+	+	+	+	No	No	3	Crop is harvested prior to bloom when not grown for seed production.	
Chrysanthemum (Asteraceae)	Leafy Vegetables	+	+	+	+	No	No	81, extrapolation from pyrethrum in reference 3	Crop is harvested prior to bloom when not grown for seed production.	
Corn salad (Valerianaceae)	Leafy Vegetables	<i>Uncertainty^a</i>								Crop may be inherently attractive to bee pollinators, but harvested prior to bloom
Cress, garden (Brassicaceae)	Leafy Vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Crop is harvested prior to bloom when not grown for seed production.	
Cress, upland (Brassicaceae)	Leafy Vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Crop is harvested prior to bloom when not grown for seed production.	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes	
Dandelion (Asteraceae)	Leafy Vegetables	++	++	++	++	No	No	80, Attractiveness extrapolated from lettuce in Table 1	Harvested prior to bloom. Flowers are removed by mechanical means when not grown for seed production. Important sources of nectar and pollen for all bee species early in the spring when few other flowers are blooming. All bumble bee species use it as a food source for early brood production.	
Dock/sorrel (Polygonaceae)	Leafy Vegetables	<i>Uncertainty^a</i>								Crop may be inherently attractive to bee pollinators, but harvested prior to bloom
Endive (Asteraceae)	Leafy Vegetables	+	+	+	+	No	No	3, attractiveness extrapolated from lettuce in Table 1	Crop is harvested prior to bloom when not grown for seed production.	
Fennel (Apiaceae)	Leafy Vegetables	++	++	+	+	Yes	No	3, Attractiveness to wild bees extrapolated from chervil above	Crop is harvested prior to bloom when not grown for seed production.	
Orach (Chenopodiaceae)	Leafy Vegetables	<i>Uncertainty^a</i>								Crop may be inherently attractive to bee pollinators, but harvested prior to bloom
Parsley (Apiaceae)	Leafy Vegetables	+	+	+	+	No	No	3, attractiveness ratings extrapolated from chervil above	Crop is harvested prior to bloom when not grown for seed production.	
Purslane, garden (Apiaceae)	Leafy Vegetables	+	+	+	+	No	No	Extrapolated from chervil above	Crop is harvested prior to bloom when not grown for seed production.	
Winter purslane (Portulacaceae)	Leafy Vegetables	<i>Uncertainty^a</i>								Crop may be inherently attractive to bee pollinators, but harvested prior to bloom
Radicchio (Asteraceae)	Leafy Vegetables	+	+	N/AV	+ <i>Andrena, Anthidium, Halictus, Osmia</i>	Yes	N/AV	Attractiveness extrapolated from chicory in Table 1	Crop is harvested prior to bloom when not grown for seed production.	
Rhubarb (Polygonaceae)	Leafy Vegetables	Open pollinated, rarely self-pollinated								Crop may be inherently attractive to bee pollinators, but harvested prior to bloom

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
New Zealand spinach (Aizoaceae)	Leafy Vegetables	<i>Uncertainty^a</i>							Crop may be inherently attractive to bee pollinators, but harvested prior to bloom
Swiss chard (Chenopodiaceae)	Leafy Vegetables	-	+	N/AV	+	Yes	No	Extrapolated from sugar beet in Table 1	Crop is harvested prior to bloom when not grown for seed production. Requires pollination only for breeding; small % of acreage
Vine spinach (Basellaceae)	Leafy Vegetables	<i>Uncertainty^a</i>							Crop may be inherently attractive to bee pollinators, but harvested prior to bloom
Brussels sprouts (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Cavalo broccolo (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Collards (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Kale (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Kohlrabi (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Mizuna (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Mustard greens (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Mustard spinach (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Rape greens (Brassicaceae)	Brassica leafy vegetables	++	++	+	+	No	No	Extrapolated from mustard seed and cabbage on Table 1	Harvested prior to bloom.
Bean (lupinus) (Fabaceae)	Legume vegetable	++	+	+	+	No	No	1, 3, 91	
Bean (vigna) (Fabaceae)	Legume vegetable	+	+	+	+	No	No	1, 3	
Guar (Fabaceae)	Legume vegetable	+	+	+	+	No	No	Extrapolated from Bean (lupinus) above	
Jackbean (Fabaceae)	Legume vegetable	+	+	+	+	No	No	Extrapolated from Bean (lupinus) above	
Lablab bean (Fabaceae)	Legume vegetable	+	+	+	+	No	No	Extrapolated from Bean (lupinus) above	
Pigeon pea (Fabaceae)	Legume vegetable	+	+	+	+	No	No	Extrapolated from Bean (lupinus) above	
Sword bean (Fabaceae)	Legume vegetable	+	+	+	+	No	No	Extrapolated from Bean (lupinus) above	
African eggplant (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Bush tomato (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	Tomatoes grown in green houses require pollination by managed bumble bees.
Cocona (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Currant tomato (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Garden huckleberry (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Goji berry (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Groundcherry (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Martynia (Pedaliaceae)	Fruiting vegetable	<i>Uncertainty^a</i>							
Naranjilla (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Pea eggplant (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Pepino (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Bell pepper (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Roselle (Malvaceae)	Fruiting vegetable	+	+	N/AV	N/AV	Yes	No	71	
Scarlet eggplant (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Sunberry (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Tomatillo (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Tree tomato (Solanaceae)	Fruiting vegetable	-	-	++	+	No	No	Extrapolated from entry for eggplant in Table 1	
Chayote (Cucurbitaceae)	Cucurbit vegetable	+	+	+	+	No	No	3	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Citron melon (Cucurbitaceae) = watermelon	Cucurbit vegetable	+	+	+	+	Yes	Yes	3	
Momordica spp. (Cucurbitaceae)	Cucurbit vegetable	+	+	+	+	Yes	Yes	Extrapolated from entry above	
Calamondin (Rutaceae)	Citrus fruit	++	++	+	+	No	No	9	
Citron (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Citrus hybrids (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Kumquat (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Mediterranean mandarin (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Mount white lime (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
New guinea wild lime (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Tangelo (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Tangor (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Uniq fruit (Rutaceae)	Citrus fruit	++	++	+	+	No	No	Extrapolated from entry above	
Azarole (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Crabapple (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	Yes	95, Extrapolated from apple in Table 1 ;	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Loquat (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Mayhaw (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Medlar (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Asian pear (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	Yes	Extrapolated from apple in Table 1	
Pseudocydonia sinensis (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Tejocote (Rosaceae)	Pome fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Capulin (Rosaceae)	Stone fruit	++	+	+	++ <i>Andrena, Anthidium, Halictus, Osmia, Anthophora, Habropoda</i>	Yes	No	Extrapolated from apple in Table 1	
Jujube (Rhamnaceae)	Stone fruit	++	+	+	+	Yes	No	3,5	
Nectarine (Rosaceae)	Stone fruit	++	+	+	+	Yes	Yes	3,5	
Peach (Rosaceae)	Stone fruit	++	+	+	+	Yes	Yes	3,5	
Plum (various) (Rosaceae)	Stone fruit	++	+	+	+	Yes	Yes	3,5	
Plumcot (Rosaceae)	Stone fruit	++	+	+	+	Yes	No	Extrapolated from entry for Plum	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Sloe (Rosaceae)	Stone fruit	++	+	+	+	Yes	No	Extrapolated from entry for Plum	
Aronia berry (Rosaceae)	Berry and small fruit	+	+	+	+	Yes	No	8	
Bayberry (Myricaceae)	Berry and small fruit	<i>Uncertainty^a</i>							
Bearberry (Ericaceae)	Berry and small fruit	+	+	++	++ <i>Andrena, Colletes, Osmia, Anthophora, Xylocopa</i>	Yes	No	Extrapolated from blueberry in Table 1 , similar flower	
Bilberry (Ericaceae)	Berry and small fruit	+	+	++	++ <i>Andrena, Colletes, Osmia, Anthophora, Xylocopa</i>	Yes	No	Extrapolated from blueberry in Table 1 , similar flower	
Blackberry (Rosaceae)	Berry and small fruit	+	+	++	++	Yes	Yes	1,3,5	
Buffaloberry (Elaeagnaceae)	Berry and small fruit	<i>Uncertainty^a</i>							
Che (Moraceae)	Berry and small fruit	<i>Uncertainty^a</i>							
Chokecherry (Rosaceae)	Berry and small fruit	+	+	N/AV	N/AV	Yes	No	58, 59	
Cloudberry (Rosaceae)	Berry and small fruit	+	+	+	+	Yes	No	60, 61	
European barberry (Berberidaceae)	Berry and small fruit	<i>Uncertainty^a</i>							
Highbush cranberry (Caprifoliaceae)	Berry and small fruit	<i>Uncertainty^a</i>							
Edible honeysuckle (Caprifoliaceae)	Berry and small fruit	+	+	+	+	Yes	No	62	
Huckleberry (Ericaceae)	Berry and small fruit	+	+	++	++ <i>Andrena, Colletes, Osmia, Anthophora, Xylocopa</i>	Yes	No	Extrapolated from blueberry in Table 1 , similar flower	
Jostaberry (Grossulariaceae)	Berry and small fruit	-	+	+	+	Yes	No	63, Extrapolated from Currants in Table 1	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Juneberry (Rosaceae)	Berry and small fruit	+	+	+	+	Yes	No	7, 64	
Lingonberry (Ericaceae)	Berry and small fruit	+	+	+	+	Yes	No	65, 66	
Maypop (Passifloraceae)	Berry and small fruit	-	-	-	+ <i>Xylocopa</i>	Yes	No	67	
Mulberry (Moraceae)	Berry and small fruit	-	-	-	-	No	No	68	Wind pollinated
Partridgeberry (Rubiaceae)	Berry and small fruit	-	-	+	-	Yes	No	69	
Phalsa (Tiliaceae)	Berry and small fruit	+	+	+	+	Yes	No	3	
Pin cherry (Rosaceae)	Berry and small fruit	++	+	+	++ <i>Osmia</i>	Yes	No		Extrapolated from cherry tree
Salal (Ericaceae)	Berry and small fruit	+	+	++	++ <i>Andrena, Colletes, Osmia, Anthophora, Xylocopa</i>	Yes	No		Extrapolated from blueberry in Table 1 , similar flower
Schisandra berry (Schisandraceae)	Berry and small fruit	<i>Uncertainty^a</i>							
Beechnut (Fagaceae)	Tree nut	-	-	-	-	No	No	70	Wind pollinated
Brazil nut (Lecythidaceae)	Tree nut	+	+	++	+	No	No	15	
Bur oak (Fagaceae)	Tree nut	+	-	-	-	No	No	16	Wind pollinated
Butternut (Juglandaceae)	Tree nut	+	-	-	-	No	No	17	
Cashew (PR)(Anacardiaceae)	Tree nut	+	+	N/AV	N/AV	Yes	No	3	
Candlenut (Euphorbiaceae)	Tree nut	<i>Uncertainty^a</i>							
Chinquapin (Fagaceae)	Tree nut	++	+	+	+	No	No	3	
Coconut (Arecaceae)	Tree nut	+	+	+	+	Yes	No	3	
Ginkgo (Ginkgoaceae)	Tree nut	-	-	-	-	No	No	18	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Guiana chestnut (PR) (Bombacaceae)	Tree nut	<i>Uncertainty^a</i>							
Heartnut (Juglandaceae)	Tree nut	+	-	-	-	No	No	Similar to butternut (above), information transferred from above	
Hickory (Juglandaceae)	Tree nut	<i>Uncertainty^a</i>							
Macadamia nut (PR) (Proteaceae)	Tree nut	+	+	N/AV	N/AV	Yes	No	3	
Pachira (Bombacaceae)	Tree nut	<i>Uncertainty^a</i>							
Peach palm nut (Arecaceae)	Tree nut	-	-	-	-	No	No	19	Pollinated by beetles
Pecan (Juglandaceae)	Tree nut	-	-	-	-	No	No	20	Wind pollinated
Pine nut (Pinaceae)	Tree nut	<i>Uncertainty^a</i>							
Tropical almond (Combretaceae)	Tree nut	+	+	N/AV	N/AV	No	No	21	
Millet (Poaceae)	Cereal grains	+	-	-	-	No	No	Similar to Grasses (Poa) so information transferred from Table 1	Source of pollen only when no other forage sources are available
Popcorn (Poaceae)	Cereal grains	+	-	-	-	No	No	Similar to Grasses (Poa) so information transferred from Table 1	Source of pollen only when no other forage sources are available
Teosinte (Poaceae)	Cereal grains	+	-	-	-	No	No	Similar to Grasses (Poa) so information transferred from Table 1	Source of pollen only when no other forage sources are available
Wild rice (Poaceae)	Cereal grains	+	-	-	-	No	No	Similar to Grasses (Poa) so information transferred from Table 1	Source of pollen only when no other forage sources are available

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Velvet bean (Fabaceae)	nongrass animal feeds	<i>Uncertainty^a</i>							
Lupin (Fabaceae)	nongrass animal feeds	+	+	+	+	No	No	5	Requires bee pollination only when grown for seed; honey bees used when grown for seed; Small % of acreage.
Crown vetch (Fabaceae)	nongrass animal feeds	+	+	++	++ <i>Megachile, Osmia</i>	Yes	Yes	Extrapolated from entry below	Requires bee pollination only when grown for seed; Small % of acreage. It is a poor seed producer as it produces little pollen or nectar,
Vetch (Fabaceae)	nongrass animal feeds	+	+	++	+ <i>Megachile, Osmia</i>	Yes	Yes	89	Requires bee pollination only when grown for seed; honey bees used when grown for seed; Small % of acreage
Milk vetch (Fabaceae) <i>Astragalus</i> spp.	nongrass animal feeds	+	+	++	+ <i>Megachile</i>	Yes	Yes	96, 97	Requires bee pollination only when grown for seed; Small % of acreage; bumble bees more effective pollinators than honey bees or leafcutter bees.
Angelica (Apiaceae)	Herbs and spices	<i>Uncertainty^a</i>							
Annatto (Bixaceae)	Herbs and spices	+	-	N/AV	N/AV	No	No	23	
Lemon balm (Lamiaceae)	Herbs and spices	+	+	+	+	No	No	24	
Basil (Lamiaceae)	Herbs and spices	+	+	+	+	No	No	7	Requires bee pollination only when grown for seed. Small % of acreage
Borage (Boraginaceae)	Herbs and spices	+	++	+	+	Yes	Yes	25, 98, 99	Requires bee pollination only when grown for seed. Small % of acreage
Burnet (Rosaceae)	Herbs and spices	+	+	+	+	No	No	26	
Camomille (Asteraceae)	Herbs and spices	+	+	N/AV	+	No	No	27, Extrapolated to potential US experience	
Black caraway (Ranunculaceae)	Herbs and spices	+	+	+	+	No	No	28, Extrapolated to potential US experience	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Caper buds (Capparaceae)	Herbs and spices	+	++	N/AV	+	No	No	29, Extrapolated to potential US experience	Requires bee pollination only when grown for seed. Small % of acreage
Catnip (Lamiaceae)	Herbs and spices	+	++	++	++	No	No	7, 30	Requires bee pollination only when grown for seed. Small % of acreage
Celery seed (Apiaceae)	Herbs and spices	+	+	+	+	Yes	No	3	Requires bee pollination only when grown for seed. Small % of acreage
Chinese chives (Liliaceae)	Herbs and spices	+	++	+	+	No	No	Extrapolated from chive, 3	Requires bee pollination only when grown for seed. Small % of acreage
Cinnamon (Lauraceae)	Herbs and spices	+	+	N/AV	N/AV	Yes	No	31	Requires bee pollination only when grown for seed. Small % of acreage
Clary (Lamiaceae)	Herbs and spices	+	+	+	+	Yes	No	7	Requires bee pollination only when grown for seed. Small % of acreage
Costmary (Asteraceae)	Herbs and spices	+	+	+	+	Yes	No	Extrapolated from chamomile, 27	Requires bee pollination only when grown for seed. Small % of acreage
Culantro (Apiaceae)	Herbs and spices	<i>Uncertainty^a</i>							
Horehound (Lamiaceae)	Herbs and spices	+	+	+	+	Yes	No	Extrapolated from 24	Requires bee pollination only when grown for seed; Small % of acreage
Hyssop (Lamiaceae)	Herbs and spices	+	+	++	++	Yes	No	7	Requires bee pollination only when grown for seed; Small % of acreage
Lavendar (Lamiaceae)	Herbs and spices	+	++	++	++	Yes	No	3, 5	Requires bee pollination only when grown for seed; Small % of acreage
Lemongrass (Graminae)	Herbs and spices	-	-	-	-	No	No	3	As a grass, primarily wind pollinated
Lovage (Apiaceae)	Herbs and spices	<i>Uncertainty^a</i>							
Mace (Myristicaceae)	Herbs and spices	-	-	-	-	No	No	32	
Marigold (Asteraceae)	Herbs and spices	+	+	-	+	No	No	55	
Marjoram (Lamiaceae)	Herbs and spices	+	+	+	+	No	No	56	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Nasturtium (Tropaeolaceae)	Herbs and spices	+	+	++	+	No	No	7	
Nutmeg (Myristicaceae)	Herbs and spices	-	-	-	-	No	No	32	
Parsley (Apiaceae)	Herbs and spices	+	+	+	+	No	No	33	Bees important for seed production
Rue (Rutaceae)	Herbs and spices	+	+	+	+	Yes	No	85	Bees important for seed production
Rosemary (Lamiaceae)	Herbs and spices	++	++	+	+	No	No	34	Perennial shrub, propagated vegetatively
Sage (Lamiaceae)	Herbs and spices	<i>Uncertainty^a</i>							
Savory (Lamiaceae)	Herbs and spices	+	+	+	+	Yes	No	86	Bees important for seed production
Tansy (Asteraceae)	Herbs and spices	+	+	+	+	No	No	35	
Tarragon (Asteraceae)	Herbs and spices	+	+	+	+	No	No	87	
Vanilla (Orchidaceae)	Herbs and spices	+	+	+	+	No	No	3	Flowers main pollinated by hand
Wintergreen (Ericaceae)	Herbs and spices	-	-	+	-	No	No	36	Flowers visited mostly by bumble bees
Wormwood (Asteraceae)	Herbs and spices	<i>Uncertainty^a</i>							
Woodruff (Rubiaceae)	Herbs and spices	<i>Uncertainty^a</i>							
Borage (Boraginaceae)	Oilseed	++	+	+	+	Yes	No	3, 37	
Calendula (Asteraceae)	Oilseed	+	+	+	+	Yes	No	3	
Chinese tallow (Euphorbiaceae)	Oilseed	+	+	+	+	Yes	No	3	
Crambe (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	3	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Cuphea (Lythraceae)	Oilseed	+	+	+	+	Yes	No	3, 10	
Echium (Boraginaceae)	Oilseed	+	+	+	+	Yes	No	3	
Euphorbia (Euphorbiaceae)	Oilseed	+	+	+	+	Yes	No	3	
Evening primrose (Onagraceae)	Oilseed	+	+	+	+	Yes	No	3	
Flax seed (Linaceae)	Oilseed	+	+	+	+	No	No	3	
Gold of pleasure (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	3	
Hare's ear mustard (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	3	
Jojoba (Simmondsiaceae)	Oilseed	+	+	+	+	Yes	No	3	
Lesquerella (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	100, 101	
Lunaria (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	3	
Meadowfoam (Limnanthaceae)	Oilseed	+	+	+	+ <i>Osmia</i>	Yes	No	3	
Milkweed (Asclepiadaceae)	Oilseed	+	+	+	+	Yes	No	88	
Niger seed (Asteraceae)	Oilseed	+	+	+	+	Yes	No	3	
Oil radish (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	3	
Rose hip (Rosaceae)	Oilseed	+	+	+	+	Yes	No	3	
Stokes aster (Asteraceae)	Oilseed	+	+	+	+	Yes	No	3	
Stokes aster (sweet rocket) (Brassicaceae)	Oilseed	+	+	+	+	Yes	No	3	
Tallowwood (Olacaceae)	Oilseed	+	+	+	+	Yes	No	3	

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Crop	EPA Crop Group	HB Poll ¹	HB Nec ¹	Bumble Bees	Solitary Bees	Requires Bee Pollination	Uses Managed Pollinators	Reference Number	Notes
Veronia (Asteraceae)	Oilseed	+	+	+	+	Yes	No	3	
Tea oil plant (Theaceae)	Oilseed	+	+	+	+	Yes	No	3	

^aWhere no data are identified for a given crop, there is uncertainty regarding its attractiveness to pollinating bees, and “*Uncertainty*” is listed in the row

¹ HB= honey bee; Poll = Pollen; Nec = Nectar

Table 3. List of references cited in Tables 1 and 2

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

Reference No.	Reference Title
1	Delaplane and Mayer 2000. Crop pollination by bees.
2	El-Wahab <i>et al.</i> 2011. Insect pollinators of anise plants (<i>Pimpinella anisum</i> L.) and the important role of honey bees (<i>apis mellifera</i> L.) on their yield productivity. Archives Of Phytopathology And Plant Protection 45
3	McGregor 1976. Insect pollination of cultivated crop plants. USDA ARS.
4	Smith <i>et al.</i> 2008. Artichoke production in California
5	Free, J. B. 1993. Insect Pollination of Crops. Academic Press, London
6	Charlebois, D., Byers, P., Finn, C.E., Thomas, A. 2010. Elderberry: Botany, Horticulture, Potential. Horticultural Reviews 37, 213-280
7	R. Isaacs, pers. obs.
8	Bader, K. L., and Anderson, S. R. (1962). Effect of pollen and nectar collecting honeybees on the seed yield of birdsfoot trefoil, <i>Lotus corniculatus</i> L. Crop Science, 2, 148-149.
9	Sanford, M.T. Pollination of citrus by honey bees. http://edis.ifas.ufl.edu/aa092
10	Parker, F.D. and Tepedino, V.J. (1990) Bee pollination of <i>Cuphea</i> (Lythraceae) species in greenhouse and field. Pan-Pacific Entomologist 66, 9-12.
11	Fohou <i>et al.</i> 2009. Pollination and yield responses of cowpea (<i>Vigna unguiculata</i> L. Walp.) to the foraging activity of <i>Apis mellifera adansonii</i> (Hymenoptera: Apidae) at Ngaoundéré (Cameroon). African Journal of Biotechnology Vol. 8, pp. 1988-1996
12	Hirota <i>et al.</i> 2012 Relative Role of Flower Color and Scent on Pollinator Attraction: Experimental Tests using F1 and F2 Hybrids of Daylily and Nightlily PLoS ONE
13	Zych, M., Stpiczyńska, M., and Roguz, K. (2014). Pollination Biology and Breeding System of European <i>Fritillaria meleagris</i> L.(Liliaceae). <i>Reproductive Biology of Plants</i> , 147.
14	Voss <i>et al.</i> Onion seed production in California. ANR publication 8008 University of California Online
15	Malhotra, SP (2008) World Edible Nuts Economy. Concept Publishing
16	Knapp, E. E., Goedde, M. A., and Rice, K. J. (2001). Pollen-limited reproduction in blue oak: implications for wind pollination in fragmented populations. <i>Oecologia</i> , 128, 48-55.
17	Burns, R.M. and Honkala, B.H. 1990. Silvics of North America. USDA Agriculture Handbook 654.
18	vanBeek, T.A. (2003) <i>Ginkgo biloba</i> . CRC Press
19	Lisbarth, C. 1996. Pollination of <i>Bactris</i> by <i>Phyllotrox</i> and <i>Epurea</i> . Implications of the Palm Breeding Beetles on Pollination at the Community Level. <i>Biotropica</i> 28, 69-81.
20	Polomski, B. (2006) Pecan planting and fertilization. http://www.clemson.edu/extension/hgic/plants/vegetables/tree_fruits_nuts/hgic1356.html
21	Thompson, L.A.J. and Evans, B. (2006) <i>Terminalia catappa</i> (tropical almond). Species profiles for pacific island agroforestry. Online at www.agroforestry.net/tti/T.catappa-tropical-almond.pdf
22	Anderson, E.J. 1959. Pollination of crown vetch. <i>Gleanings in Bee Culture</i> 87: 590 - 593.
23	Heard, T.A. (1999). THE ROLE OF STINGLESS BEES IN CROP POLLINATION. <i>Annual Review of Entomology</i> 44: 183-206.
24	Chwil, M. (2009) Flowering biology and nectary structure of <i>Melissa officianalis</i> (L.) <i>Acta Agrobotanica</i> 62: 23-30.
25	Carreck, N.L. and Williams, I.H. (2002). Food for insect pollinators on farmland: insect visits to flowers of annual seed mixtures. <i>Journal of Insect Conservation</i> 6, 13-23.
26	Anon. Plant Guide for Small Burnet (<i>Sanguisorba minor</i>) USDA NRCS Plant Guide. Online at: plants.usda.gov/plantguide/pdf/pg_sami3.pdf
27	Kuberappa, G. C.; Shilpa, P.; Vishwas, A. B.; Vasundhara, M. (2008) Insect pollinator fauna, abundance and their foraging activity on chamomile (<i>Matricaria chamomilla</i> L.). <i>Biomed</i> 2, 345-349
28	Abd El-Wahab, T.E. and Ebadah, I.M.A. (2011) Impact of honeybee and other insect pollinators on the seed setting and yield production of black cumin <i>Nigella sativa</i> L. <i>J. Basic. Appl. Sci. Res.</i> , 1 622-626.
29	A Dafni, D Eisikowitch, Y Ivri (1987) Nectar flow and pollinators' efficiency in two co-occurring species of <i>Capparis</i> (Capparaceae) in Israel. <i>Plant Systematics and Evolution</i> 157, 181-186
30	Sih, A. and Baltus, M.-S. (1987) Patch Size, Pollinator Behavior, and Pollinator Limitation in Catnip. <i>Ecology</i> 68:1679-1690
31	Ravindran, P.N., Nirmal-Babu, K. and Shylaja, M. (2003) <i>Cinnamon and Cassia: The Genus Cinnamomum</i> . CRC Press.
32	Sharma, M.V. and Armstrong, J.E. (2013) Pollination of <i>Myristica</i> and other nutmegs in natural populations. <i>Tropical Conservation Science – Special Issue Vol.6:</i> 595-607.
33	Burgett, M. (1980) Pollination of parsley (<i>Petroselinum crispum</i>) grown for seed. <i>Journal of Apicultural Research</i> 19, 79-82.

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

34	Keasar, T., Sadeh, A., and Shmida, A. (2001) Variability in nectar production and standing crop, and their relation to pollinator visits in a Mediterranean shrub. <i>Arthropod-Plant Interactions</i> 2, 117-123.
35	Gucker, Corey L. 2009. <i>Tanacetum vulgare</i> . In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: http://www.fs.fed.us/database/feis/ [2014, February 9].
36	Mirick, S. and Quinn, J.A. (1981) Some observations on the reproductive biology of <i>Gaultheria procumbens</i> (Ericaceae). <i>Amer. J. Bot.</i> 68,
37	Davis, A.R., Mitchell, S. and Junor, D. The Importance of Honey Bees (<i>Apis mellifera</i> L.) as Pollinators of OF Borage (<i>Borago officinalis</i> L.) in Saskatchewan. Online at http://www.usask.ca/soilscrops/conference-proceedings/previous_years/Files/97/1997docs/364.pdf
38	Kays, SJ and Nottingham, SF (2008) <i>Biology and Chemistry of Jerusalem Artichoke: Helianthus tuberosus</i> L. CRC Press, Boca Raton, FL
39	Anon. Peppermint. http://oregonstate.edu/dept/coarc/peppermint-0
40	Sanchez C. (2010) Cultivation of <i>Pleurotus ostreatus</i> and other edible mushrooms. <i>Applied Microbiology and Biotechnology</i> 85: 1321-1337.
41	Wolfe, J.A. (1992) <i>Sweetpotato, an untapped food resource</i> . Cambridge University Press.
42	Anon. <i>Vicia sativa</i> Linnaeus. Plant Diversity Website. University of Michigan. climbers.lsa.umich.edu/wp-content/uploads/.../ViciasatiFABAFINAL.pdf
43	Cervenková Z and Münzbergová Z. (2014) Pollen limitation and pollinator preferences in <i>Scorzonera hispanica</i> . <i>Plant Biol (Stuttg)</i> . doi: 10.1111/plb.12142. [Epub ahead of print]
44	Expert opinion (H. Burrack, NC State University Extension Specialist in Tobacco) and see http://www.tobacco.ncsu.edu/files/2013%20Flue-Cured%20Guide.pdf
45	UC Davis - Walnuts in California. http://fruitandnuteducation.ucdavis.edu/education/fruitnutproduction/Walnut/
46	Produced vegetatively from tubers. http://www.extento.hawaii.edu/kbase/reports/taro_prod.htm
47	Parsnip production guide, Oregon State University. http://nwrec.hort.oregonstate.edu/parsnip.htm
48	Sources: USDA, National Agricultural Statistics Service, Crop Production and Crop Values.
49	Arista, M., Ortiz, P. L., and Talavera, S. (1999). Apical pattern of fruit production in the racemes of <i>Ceratonia siliqua</i> (Leguminosae: Caesalpinioideae): role of pollinators. <i>American journal of botany</i> , 86(12), 1708-1716.
50	Olsen, J. L., Mehlenbacher, S. A., and Azarenko, A. N. (2000). Hazelnut pollination. <i>HortTechnology</i> , 10(1), 113-115.
51	Small, E., and Antle, T. (2003). A preliminary study of pollen dispersal in <i>Cannabis sativa</i> in relation to wind direction. <i>Journal of Industrial Hemp</i> , 8(2), 37-50.
52	F.J. Muehlbauer, R.J. Summerfield, W.J. Kaiser, S.L. Clement, C.M. Boerboom, M.M. Welsh-Maddux, and R.W. Short Principles and Practice of Legume Production. Online at http://www.ars.usda.gov/is/np/lentils/lentils.htm#Background
53	Herrera, E. 1997 Growing pistachios in New Mexico. Cooperative Extension Service Circular 532, New Mexico State University. http://aces.nmsu.edu/pubs/_circulars/CR532.pdf
54	Free, J. B., and Nuttall, P. M. (1968). The pollination of oilseed rape (<i>Brassica napus</i>) and the behaviour of bees on the crop. <i>The Journal of Agricultural Science</i> , 71(01), 91-94.
55	Bhardwaj, H., Thaker, P., and Srivastava, M. (2010). Hymenopteran visitors of <i>Tagetes erecta</i> as observed in an agro-ecosystem near Bikaner, Rajasthan. <i>Current Biotica</i> , 4(1), 94-102.
56	Comba, L., Corbet, S. A., Hunt, L., and Warren, B. (1999). Flowers, nectar and insect visits: Evaluating British plant species for pollinator-friendly gardens. <i>Annals of Botany</i> , 83(4), 369-383.
57	http://extension.umass.edu/floriculture/fact-sheets/production-hybrid-lilies-pot-plants
58	Danka, R. G., and Beaman, L. D. (2007). Flight activity of USDA-ARS Russian honey bees (Hymenoptera: Apidae) during pollination of lowbush blueberries in Maine. <i>Journal of economic entomology</i> , 100(2), 267-272.
59	Shiell, K. J., St-Pierre, R. G., and Zatylny, A. M. (2002). Timing, magnitude and causes of flower and immature fruit loss in pin cherry and choke cherry. <i>Canadian journal of plant science</i> , 82(1), 157-164.
60	Brown, A. O., and McNeil, J. N. (2009). Pollination ecology of the high latitude, dioecious cloudberry (<i>Rubus chamaemorus</i> ; Rosaceae). <i>American journal of botany</i> , 96(6), 1096-1107.
61	Pelletier, L., Brown, A., Otrysko, B., and McNeil, J. N. (2001). Entomophily of the cloudberry (<i>Rubus chamaemorus</i>). <i>Entomologia Experimentalis et Applicata</i> , 101(3), 219-224.
62	Bors, B. Haskap Pollination Strategy - http://www.fruit.usask.ca/articles/pollinationstrategy.pdf
63	Barney, D.L. and Fallari, E. (2009) Currants, gooseberries, and jostaberries in the inland northwest and intermountain west. University of Idaho Extension Bul 855.
64	Gorchov, D. L. (1988). Effects of pollen and resources on seed number and other fitness components in <i>Amelanchier arborea</i> (Rosaceae: Maloideae). <i>American Journal of Botany</i> , 1275-1285.
65	Holloway, P.S. (1984) Lingonberry cultivation. <i>Agroborealis</i> , July issue, 15-20.

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

66	Holloway, P. S., Kruse, J. J., and Davis, A. N. (2002, August). Insect visitors and potential pollinators of lingonberries, <i>Vaccinium vitis-idaea</i> subsp. minus, in sub-arctic Alaska. In <i>XXVI International Horticultural Congress: Berry Crop Breeding, Production and Utilization for a New Century 626</i> (pp. 433-438).
67	McGuire, C. M. (1998). Field performance and phenotypic variation of <i>Passiflora incarnata</i> L. in New York State. <i>HortScience</i> , 33(2), 240-241.
68	Janick, J and Paull, R.E. (2008) The Encyclopedia of Fruits and Nuts. CABI, Wallingford.
69	Hicks, D. J., Wyatt, R., and Meagher, T. R. (1985). Reproductive biology of distylous partridgeberry, <i>Mitchella repens</i> . <i>American Journal of Botany</i> , 1503-1514.
70	Thomas, P.A. (2000) Trees, their Natural History. Cambridge University Press, Cambridge.
71	Abdel-Moniem, A. S. H., Abd El-Wahab, T. E., and Farag, N. A. (2011). Prevailing insects in Roselle plants, <i>Hibiscus sabdariffa</i> L., and their efficiency on pollination. <i>Archives of Phytopathology and Plant Protection</i> , 44(3), 242-252.
72	Yadav, S.S. and Chen, W. (2007) Chickpea Breeding and Management. CABI, Wallingford.
73	MSU Native Plant Facts Buckwheat
74	Arista <i>et al.</i> 1999. Apical pattern of fruit production in the racemes of <i>Ceratonia siliqua</i> (Leguminosae: Caesalpinioideae): Role of Pollinators.
75	Duke 1980. Pollinators of Panax. <i>Castanea</i> 47: 261-265.
76	Walters, S.A. and E.A. Wahle. 2010. Horseradish Production in Illinois. <i>Hort Technology</i> 20 (2): 267-276.
77	Ayansola, A.A. and O. Awolowo. 2012. Honeybee Floral Resources in Southwestern Nigeria. <i>Journal of Biology and Life Science</i> . 3(1): 127-139.
78	Huaman, Z. 1992. Systematic Botany and Morphology of the Sweetpotato Plant. Technical Information Bulletin 25. International Potato Center. Lima, Peru.
79	Real, L.A. 1981. Nectar Availability and Bee-Foraging on Ipomoea (Convolvulaceae). <i>Reproductive Botany</i> . 64-69.
80	Porter, B. and C. Brenzil. 2003. Farm Facts: Dandelion Production. Saskatchewan Agriculture, Food, and Rural Revitalization.
81	USDA ARS. 2010. Plants Attractive to Native Bees: http://www.ars.usda.gov/Main/docs.htm?docid=12052
82	Lin, S., Chang, S., and Chen, S. 1993. The study of bee-collected pollen loads in Nantou, Taiwan. <i>Taiwania</i> 38:117-133.
83	Schmidt, M.R. and Bothma, G. Indications of Bee Pollination in Sorghum and its implications in transgenic biosafety. <i>ISMN</i> . 46.
84	Ippolito, A., Fernandes, G.W., and Holtsford, T.P. Pollinator preferences and relative fitness of <i>Nicotiana glauca</i> , <i>N. glauca</i> and their F1 hybrids.
85	Ren, M., Tang, J. 2012. Up and down: stamen movements in <i>Ruta graveolens</i> (Rutaceae) enhance both outcrossing and delayed selfing. <i>Annals of Botany</i> .
86	Kargar, V., Alizadeh, A., and Namayandeh, A. 2014. Essential oil constituents of <i>Satureja sahendica</i> Bornm. And <i>Satureja hortensis</i> L. cultivated in Iran. <i>International Journal of Farming and Allied Sciences</i> .
87	Makkdoom Sabir, A. 2011. Diversity of <i>Bombus</i> species (Apidae: Hymenoptera) and utilization of food resources in Northern Pakistan. University of Agriculture, Faisalabad Pakistan. Thesis.
88	Willson, M.F. and R. I. Bertin. 1979. Flower-visitors, nectar production, and inflorescence size of <i>Asclepias syriaca</i> . <i>Canadian Journal of Botany</i> , 57(12): 1380-1388
89	Richards KW. 1997. The alfalfa leafcutter bee as a potential pollinator of some annual clover species. <i>J Apiculture Res.</i> 34(3):115-121
90	Williams I.H. 1988. The pollination of linseed and flax. <i>Bee World</i> . 69:149-152
91	Williams I.H., A. P. Martens, A.W. Ferguson, and S.J. Clark. 1990. Effect of pollination on flower, pod and seed production in white lupin (<i>Lupinus albus</i>). <i>J Agric Sci, Cambridge</i> 115:67-73
92	JAC Miller, L Henning, V.L. Heazlewood, P.J. Larkin, J. Chitty, R. Allen, P.H. Brown, W. Gerlach, and A. J. Fist. 2005. Pollination biology of oilseed poppy, <i>Papaver somniferum</i> L. <i>Australian Journal of Agricultural Research</i> 56(5) 483–490.
93	A. K. Pandey and A. Kumari. Pollination ecology of safflower (<i>Carthamus tinctorius</i> linn). Proceeding 7 th International safflower conference, Australia 10g www.australianoilseeds.com/ data/./final_Pandey_poster_paper.pdf
94	Zheleznov AV, N. B Zheleznova, and V. K. Shumnyi VK. 2001. Seed set analysis in three amaranth species (<i>A. cruentus</i> , <i>A. caudatus</i> , and <i>A. lividus</i>) under self- and cross-pollination conditions. <i>Tsitol Genet.</i> 35(1):39-45
95	2013 Crabapple Pollenizers for Apples – Ontario www.omafra.gov.on.ca/english/crops/facts/00-011.htm/
96	Richards KW. 1986. Pollination requirements of cicer milkvetch, <i>Astragalus cicer</i> . <i>Journal of Range Management</i> 39(5):457-459.
97	Richards KW. 1987. Diversity, density, efficiency and effectiveness of pollinators of cicer milkvetch, <i>Astragalus cicer</i> L. <i>Can J Zoology</i> 65(9): 2168-2176.

Attractiveness of Agricultural Crops to Pollinating Bees for the Collection of Nectar and/or Pollen, 2015

98	Pollination of Crops - Welcome to the Red River Apiarists www.beekeepingmanitoba.com/main
99	Davis A <i>et al.</i> 1997. The importance of honey bees (<i>Apis mellifera</i> L.) as pollinators of Borage (<i>Borago officinalis</i> L.) in Saskatchewan http://www.usask.ca/soilscrops/conferenceproceedings/previous_years/Files/97/1997docs/364.pdf
100	R. J. Mitchell. 1997. Effects of Pollination Intensity on <i>Lesquerella fendleri</i> Seed Set <i>Oecologia</i> 109 (3):382-388
101	R. J. Mitchell. 1995 Effects of Pollination Method on Paternal Success in <i>Lesquerella fendleri</i> . <i>American Journal of Botany</i> 82(4): pp. 462-467.
102	Richards, K.W., Edwards, P. 1988. Density, diversity and efficiency of pollinators of <i>Onobrychis viciaefolia</i> . <i>Can Ent</i> 120: 1085- 1100)
103	Richards, K.W., 2003. Potential use of the alfalfa leafcutter bee <i>Megachile rotundata</i> to pollinate sweet clover. <i>J. Apicultural Res.</i> 42(1-2): 21-24.
104	Moffett, J.O., L.S. Stith, C.C. Burkhart and C.W. Shipman. 1975. Honey Bee Visits to Cotton Flowers, <i>Environ. Entom.</i> 4(2): 203-206.
105	McGregor, S.E. 1959. Cotton-Flower Visitation and Pollen Distribution by Honey Bees, <i>Science</i> 9, January, 129(3341): 97-98.
106	Meredith, W.R., Jr. and R.R. Bridge. 1973. Natural Crossing in Cotton (<i>Gossypium hirsutum</i> L.) in the Delta of Mississippi, <i>Crop Science</i> , Sept.-Oct., 13: 551-552.
107	Vansell, G.H. 1944. Cotton Nectar in Relation to Bee Activity and Honey Production. <i>J. Econ. Entomol.</i> , 37(4): 528-530.
108	A. Whalen, A. Catchot, J. Gore, S. Stewart, G. Lorenz, D. Cook, and F. Musser. 2015. In, <i>Proc. Beltwide Cotton Production Conferences</i> .
109	Vaissière, B.E. and S.B. Vinson, 1994. Pollen morphology and its effect on pollen collection by honey bees, <i>Apis Mellifera</i> L. (Hymenoptera: Apidae), with special Reference to Upland Cotton, <i>Gossypium Hirsutum</i> L. (Malvaceae), <i>Grana</i> 33:128-138.
110	Hutmacher, R.B., R.N. Vargas, and S.D. Wright. 2006. <i>Methods to Enable Coexistence of Diverse Cotton Production Systems</i> , University of CA ANR, pub 8191, 6pp.
111	Umbeck, P.F., K.A. Barton, E.V. Nordheim, J.C. McCarty, W.L. Parrott and J.N. Jenkins, 1991. Degree of Pollen Dispersal by Insects from a Field Test of Genetically Engineered Cotton, <i>J. Econ. Entomol.</i> 84(6): 1943-1950.